

CLASSIFICATION **CONFIDENTIAL**  
 CENTRAL INTELLIGENCE AGENCY  
 INFORMATION FROM  
 FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT

CD NO.

50X1-HUM

COUNTRY USSR  
 SUBJECT Scientific - Chemistry  
 HOW PUBLISHED Bimonthly periodical  
 WHERE PUBLISHED Moscow  
 DATE PUBLISHED May - Jun 1950  
 LANGUAGE Russian

DATE OF INFORMATION 1950

DATE DIST. 22 Aug 1950

NO. OF PAGES 2

SUPPLEMENT TO REPORT NO.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF ESPIONAGE ACT 80 U. S. C., 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED.

THIS IS UNEVALUATED INFORMATION

SOURCE Zhurnal Analiticheskoy Khimii, Vol V, No 3, 1950

QUANTITATIVE SEPARATION OF ARGON FROM SMALL QUANTITIES OF KRYPTON AND XENON BY REPEATED ADSORPTION AND DESORPTION.

E. K. Gerling  
 N. V. Baranovskaya  
 Submitted 28 Jan 1950

This research was undertaken because of the fact that although there had been numerous previous investigations (1-6) concerned with the separation of these gases from each other, none of them had considered the problem of the separation of a large quantity of argon from small quantities of the other two rare gases.

The method suggested by Peters and Well (6) was employed, and through the study of the isotherms of adsorption and desorption of argon and krypton at the three temperatures, minus 120, minus 100, and minus 80 degrees centigrade and pressures of from 1 to  $1 \cdot 10^{-4}$  millimeters, the following conclusions were obtained:

1. At these three temperatures, all of the argon which had been adsorbed by the carbon was not desorbed, and the amount not desorbed was increased as the temperature was decreased.

2. The most rapid separation of argon and krypton can be accomplished at minus 100 degrees centigrade by carrying out consecutive adsorption and desorption of the gaseous mixture over four different carbon portions, then liberating the adsorbed gases by heating, and finally repeating these operations on all carbon portions one more time. This procedure resulted in more than 99 percent of the krypton being adsorbed on the carbon, and then all of the argon was quantitatively desorbed and pumped off.

## BIBLIOGRAPHY

1. Valentiner and Schmidt, Ber. Berl. Akad., 816, 1905.
2. Dewar, Proc. Roy. Soc., XVIII, 433, 747, 1906.

**CONFIDENTIAL**

- 1 -

CLASSIFICATION		CONFIDENTIAL	
STATE	<input checked="" type="checkbox"/> NAVY	<input checked="" type="checkbox"/> NSRB	DISTRIBUTION
ARMY	<input checked="" type="checkbox"/> AIR	<input checked="" type="checkbox"/> FBI	

**CONFIDENTIAL**

CONFIDENTIAL

50X1-HUM

3. Lepape, C. R., CLXXXVII, 231, 591, 1928.
4. Moore, Proc. Roy. Soc., London, A., LXXXI, 195, 1908
5. Aston, Proc. Roy. Soc., London, A., CIII, 462, 1923.
6. Peters and K. Weil, Z. phys. Chem., A., CXLVIII, 1930; A. ang Chem., L, 43, 1930.

- E N D -

- 2 -

CONFIDENTIAL

**CONFIDENTIAL**